

CrossRef DOI of original article:

# 1 Distal Radius Fracture: A Systematic Review of Observational 2 Studies

3

4 *Received: 1 January 1970 Accepted: 1 January 1970 Published: 1 January 1970*

5

---

## 6 **Abstract**

---

8 **Index terms—**

## 9 **1 INTRODUCTION**

10 Distal Radius Fractures (DRF) are anatomically defined as those occurring within 3 cm of the radiocarpal joint,  
11 corresponding to one sixth of all fractures treated in emergency units and with great cost to the health system.  
12 1,2 The DRF have significant epidemiological and clinical-surgical importance due to the high prevalence, that  
13 is, up to 31%. 3 In addition, the complexity of the lesions varies according to the involvement of adjacent  
14 anatomical regions, implying different prognoses whose treatment may be different. conservative or surgical.  
15 [4][5][6] Observational Studies (OS) have different degrees of reliability, and can be compared to a photograph  
16 of the population (cross-sectional study) or temporal analysis of the sample in question (longitudinal study). 7  
17 Thus, the present scientific dissertation aims to explain the OS of these two methodologies about the epidemiology  
18 associated with the trauma of DRF and factors implicating the seasonality of the prevalence and incidence.

## 19 **2 II. METHODOLOGY**

20 The methodology used was the systematic review, in which the research platforms MEDLINE® and LILACS for  
21 BIREME, PUBMED and SCIELO were used. In this context, according to the Descriptors in Health Sciences  
22 platform and with descriptors in Portuguese, English and Spanish, 2 Volume 23 | Issue 5 | Compilation 1.0 the  
23 following descriptors were selected: "Bone fracture" and "Radius fracture"; "Wrist joint"; "radius fracture".

24 The inclusion criteria corresponded to complete and available articles, from the last five years (October 30,  
25 2016 the same date of the year 2021), only with human beings, OS of the transverse and longitudinal types and  
26 that addressed the distal third of the radius as an outcome.

27 Exclusion criteria were articles that did not address radius fracture, protocol validation studies, case reports  
28 and series, systematic reviews, meta-analysis, randomized controlled trials and case-control studies.

29 Table 1 summarizes the methodology, sample size, study design, primary out-come result, gender and age group  
30 included in the studies eligible for this systematic review. elimination of duplicates, 64 articles were selected for  
31 reading in full, with 18 being eligible for this systematic review. The flowchart (figure ??)  
32 demonstrates such steps of the methodological process.

33 Figure 1

34 Table 2 explains the fracture rate according to the AO 8 classification (types A, B and C) and the frequencies  
35 of accidents associated with the trauma mechanism.

## 36 **3 IV. DISCUSSION**

37 When assessing whether obesity increases the severity of DRF, in one study, it was found that there was no  
38 correlation based on a sample of 114 patients, although this population is more susceptible to fractures in this  
39 region. 9 A study aimed at quantifying the ability of unicortical distal screws to maintain the operative reduction  
40 of DRF in adults, in a sample of 75 patients undergoing volar fixation with a locking plate, resulted in data that  
41 corresponded to effective fixation and maintained the operative reductions in DRF, whereas it had the potential  
42 to decrease the incidence of extensor tendon ruptures in a 12-week post-surgical follow-up. 10 Comparing type B

### 3 IV. DISCUSSION

---

43 DRF with and without involvement of the semilunar facet treated with a volar blocking plate, we retrospectively  
44 analyzed the individuals with involvement of the semilunar facet (n=21) and the others without this involvement  
45 (n = 72), it was found that patients with DRF with involvement of the semilunar facet would have slower recovery  
46 with regard to wrist flexion, supination, ulnar deviation and greater risk of loss of both reduction and final joint  
47 step. 11 Thus, fractures of DRF with involvement of the semilunar facet present clinical factors of severity and  
48 longer time for convalescence of the bone material.

49 When studying patients aged 50 years and over with a total sample of 25,454 fractures, of which 42.1%  
50 corresponded to fractures of the radius or ulna, Ogliari et al. 12 verified that most of the injuries in this region  
51 occurred in the winter period, in which frosty days were directly associated with fractures of the radius or  
52 ulna. In addition, confirmation of the increase in accidents involving DRF in colder seasons can help direct  
53 financial resources from health services and increase the number of employees available during this period. Thus,  
54 considering the temperate climate of the study site, it can be suggested that other parts of the globe, including  
55 the southern region of Brazil, may have a similar rate of involvement, which should be corroborated by future  
56 studies and evaluated according to the possibility of greater exposure. to falls from standing height by the general  
57 population and higher risk of falls in geriatric individuals on slippery floors.

58 Another study looked at the profile of DRF in relation to climate, in which a retrospective analysis with 8,380  
59 patients verified wrist joint fractures in all women and men aged ?50 years, and demonstrated an increase in  
60 the rate of this type of fracture with the reduction in temperature (milder seasons of the year) and the average  
61 rate was 2.9 fractures/day out of a total of 2,922 days analyzed and an increase of 840 in the number of Distal  
62 Radius Fracture: A Systematic Review of Observational Studies 6 Volume 23 | Issue 5 | Compilation 1.0 6  
63 procedures during the winter period. 13 Thus, in regions with a cold climate, health services can direct resources  
64 to this patient profile. Furthermore, Range of Motion Limitation (RML) after DRF was studied in 88 geriatric  
65 patients, in whom daily rehabilitation was applied for 30 minutes for 8 weeks after the fracture, and it was  
66 verified that individuals with early stiffness (<3 months after the fracture) had greater RML compared to those  
67 with late stiffness (>3 months). 14 A cohort with a sample of 202 children (up to 16 years of age) addressed the  
68 complication rate of forearm fractures after treatment with stable intramedullary elastic nails, which correspond  
69 to in situ refracture (1.5%), refracture after nail removal (3.5%), vicious junction (1.0%), rupture of the extensor  
70 pollicis longus tendon (1.5%), infection 1 (0.5%) and reduced range of motion (1.0%). 15 Wang et al. ??6  
71 analyzed, by means of a longitudinal retrospective study, 410 children and adolescents (aged from 6 to 18 years),  
72 whose aim was to characterize polytrauma due to sports, in which it was verified that radius fractures were the  
73 most common (24.9%) relative to other long bones. In addition, males had a significantly higher rate of fractures  
74 and associated nerve injuries, with peaks of incidence in the summer and the most associated sport was basketball  
75 (28.9%).

76 A prospective longitudinal study followed patients with DRF for 12 months in order to verify the future risk  
77 of fracture in two groups, that is, young versus elderly men (65 years or older), whose result associated that the  
78 second group was more prone to having a disability in working with hands, arms and shoulder ipsilateral to the  
79 injury, regardless of the macroscopic radiographic finding. Furthermore, the decreased bone strength of these  
80 patients was associated with an increased risk of fracture at 10 years. ??7 Furthermore, another population-based  
81 retrospective longitudinal analysis sought to determine the proportion of DRF treated without adequate medical  
82 follow-up after initial care, as well as the type of medical care provided by different hospitals and physicians.  
83 And, from the analysis of 70,801 fractures, it was found that 20.8% (n=14,742) of fractures were treated without  
84 continuous medical follow-up after initial care and treatment by a small hospital emergency department, pediatric  
85 specialty or subspecialty in a pediatric emergency, were more likely to result in no follow-up. In addition, small  
86 hospitals and living in a rural area were significantly associated with non-monitoring after the injury. ??8 A  
87 retrospective observational cohort with the aim of determining the incidence of DRF, in which 90,970 DRF were  
88 identified between the years 2005 to 2012, whose incidence rate during the entire period analyzed was 52.9/10,000  
89 people/year, with the distribution between genders equal in the age group from 0 to 10 years old, however, higher  
90 in males from 11 to 17 years old. Furthermore, there was a significant variation in incidence throughout the  
91 year, with higher peaks in May (68.7/10,000 person-years) and September (73.2/10,000 person-years). ??9 Thus,  
92 considering that the period observed of higher incidences corresponds to milder temperatures, it can be suggested  
93 that such incidence may be related to sports activities or activities of greater impact.

94 A prospective cross-sectional study compared pediatric patients (<18 years) and adults (? 18 years) in two  
95 moments, before and during the covid-19 pandemic, with the aim of characterizing variation in epidemiological  
96 data on hospitalization and the need for a surgical approach. The first group showed a decrease in hospitalizations  
97 (3.8%), hospitalizations with surgical treatment (11.5%) and patients undergoing conservative treatment (7.2%).  
98 The adult population showed a decrease in the rate of hospitalizations treated surgically (12.7%) and in the  
99 number of individuals undergoing conservative treatment (30.3%), while those who underwent surgical treatment  
100 with fixation by volar plate increased substantially (275%). ??0 Rundgren et al. ??1 with the aim of determining  
101 Surgical Site Infections (SSI) after DRF surgery using different techniques (plate fixation, percutaneous pinning  
102 and external fixation), as well as factors associated with SSI in a sample of 31,807 patients, found that the rate  
103 of SSI corresponded to rates of 5%, 12% and 28%, respectively. Furthermore, it was found that the type of  
104 open fracture and being male were associated with SSI. Nagai et al. ??2 evaluated the relationship between  
105 Potentially Inappropriate Medications (PIM), activities of daily living and subsequent falls in elderly patients

106 with DRF, aged 65 years and over and divided into two groups (a group using PIM and a group not using PIM).  
107 The prevalence of prescriptions for PIM was 42.3% and their use hindered the improvement in activities of daily  
108 living and was associated with an increase in subsequent falls.

109 A retrospective cohort study included 304 adults aged 60 years or older who had isolated DRF and divided into  
110 two groups: group I with 187 participants (volar locking plate, percutaneous pinning or external fixation) and  
111 group II with 117 individuals (treated with a cast) and classified into highly and less active based on the degree of  
112 physical activity prior to the injury. The results suggested that more physical activity practiced before the injury  
113 was associated with better functional results and patient-reported self-improvement. Thus, supervised physical  
114 activity, due to the risk of falls, should be encouraged in these patients. ??3 Another research, when evaluating  
115 Parkinson's Disease (PD) in two groups with (n=23) and without (n=65) the disease regarding the best outcome  
116 in patients with DRF, in which both groups underwent open reduction followed by of internal fixation, with the  
117 aim of verifying whether the PD group would have a lower result after surgery compared to non-PD patients,  
118 it was found that there was a shorter time and a significant rate of treatment failure, these being 39.1% and  
119 4.6%, respectively. ??4 A retrospective longitudinal study with the objective of investigating the association  
120 between nutritional status and functional prognosis in elderly patients with DRF in elderly individuals, found  
121 that a positive association between malnutrition and the ability to resume activities of daily living after DRF  
122 and low levels of albumin serum levels may increase the risk of subsequent falls, and a rate of 13.5% of patients  
123 with DRF had malnutrition. ??5 When assessing the frequency with which children younger than 10 years old  
124 undergo a potentially unnecessary closed reduction associated with sedation for the DRF procedure and the cost  
125 implications, Orland et al. ??4 found that among 258 participants, 142 (55%) underwent this procedure and 38  
126 children (27%) were considered potentially unnecessary with a cost increase of about 8 times the amount and  
127 the fractures could have been treated with in situ immobilization. ??6

## 128 **4 V. FINAL CONSIDERATIONS**

129 It is verified that there is seasonality in the DRF regarding the seasons. The reduced bone strength of these  
130 patients was associated with an increased risk of fracture in 10 years, and individuals from a rural environment  
131 have less hospital support and a greater chance of not having adequate post-fracture follow-up. In addition,  
obesity was not a serious factor for DRF recovery. <sup>1</sup>



Figure 1: 4

Figure 2: Distal

Acosta Gross-

Obesity has been assessed

There was no correlation

Olivo sectional 114 to increase the severity of DRF.  
et Prospective  
al. al.

The ability of distal  
unicortical screws to

between obesity and severity of

Darda Longitudinal 75 maintain operative DRF  
et  
al. Prospective reduction in adults was  
verified.

To verify the results of the  
volar locking plate for the

Unicortical distal fixation  
during volar locking plate  
fixation effectively produced  
operative reductions in DRF.

treatment of type B DRF in-

slower recovery when compa-

Zhang Retrospective 93 involving the semilunar facet and compare with fractures red to DRF without the involve-  
et Longi-  
al. tudi-  
nal  
without this involvement.

ment to DRF without the involve-  
ment of the semilunar facet would have a  
topography.

Oglian Retrospective It aimed to explore fragility  
et  
al. Longitudinal 25 45 fractures in adults over 50

years of age, including DRF.  
To verify the incidence of

The climate can modulate the  
seasonality of fractures and,  
consequently, the use of  
health service resources.

Johns Retrospective fractures of the wrist joint in  
et

al. Longitudinal 83 80 relation to the hot and cold  
years of age, including DRF.  
To verify the incidence of

There was an increase in the

rate of this type of fracture in  
the coldest seasons of the  
year.

To verify the benefit of  
rehabilitation

regarding

Benefit of rehabilitation in

Zhang Longitudinal 88 RML due to DRF in individuals over 65 years of  
et Prospective  
al. tive

case of RML due to DRF with

| Author   | AO Classification                        | Fracture Rate According to the Trauma Mechanism   |
|--|--|---|
| Acosta-<br>Olivo<br>et<br>al.                              | A (20,18%);<br>B (32,45%);<br>C (47,37%) | Type A fractures were the most common and the most severe type of fracture, type C, was the least common in all patients (normal weight, over-weight, obese)  |
| Dardas<br>et al.   | A (40%);<br>B (12%);<br>C (48%)          | The mechanisms most associated with trauma were falls (68%) and falling from standing height (19%)  |
| Zhang<br>et al.<br>Ogliari<br>et al.<br>Johnson<br>et al.. | A (19.32);<br>B (12.50);<br>C 65.91)     | Electric bicycle accidents were the most common cause (51.6%) of all injuries, followed by falls from heights (24.7%), motor vehicle accidents  |
| Zhang<br>et al.  | (14,0%)                                  | (18.3%) and sports injuries (2.2%)  |
| Kruppa<br>et al.   |  | Drop (playing, jumping, skating, others) of 98.0%<br>Polytrauma (motor vehicle accident; fall from a height of 3 m), two accidents (1.0%) No adequate trauma (osteogenesis imperfecta; juvenile bone cyst) 2 (1.0%) |
|  |  | The most common etiologies were playing basketball (27.5%) in the male group and walking (24.0%) in the female group.   |
|  |  | The most common etiologies and locations were playing basketball (34.0%) and FR (26.2%) in the 12-15 age group, playing basketball (31.7%) and FR (23.0%) in the age 15 to 18 years                                 |
| Wang<br>et al.   |  | The most common FR fracture sites were in basketball (28.9%) and cricket (37.5%) players.   |
| Egund<br>et al.  |  |   |

---

133 [ Distal Radius Fracture: A Systematic Review of Observational Studies] , Compilation 1.0. *Distal Radius Frac-*  
134 *ture: A Systematic Review of Observational Studies* 8 (5) .

135 [Dardas et al. ()] ‘A prospective observational assessment of unicortical distal screw placement during volar plate  
136 fixation of distal radius fractures’. A Z Dardas , C A Goldfarb , M I Boyer , D A Osei , C J Dy , R P Calfee  
137 . *The Journal of hand surgery* 2018. 43 (05) p. .

138 [Amatuzzi ()] ‘Análise da evolução qualitativa de publicações em Ortopedia e Traumatologia: comparação entre  
139 a ”Revista Brasileira de Ortopedia” e’. Mll Amatuzzi . *The Journal of Bone and Joint Surgery* 2004. 39 (9)  
140 p. . (Revista Brasileira de Ortopedia)

141 [Zhang et al. ()] ‘Comparative study of type B distal radius fractures with and without lunate facet involvement  
142 treated by volar locking plate, an observational study’. X Zhang , Y Zhao , C Hu . *International Journal of  
143 Surgery* 2017. 44 p. .

144 [Acosta-Olivo et al. ()] ‘Correlation between obesity and severity of distal radius fractures’. C Acosta-Olivo , J  
145 C Gonzalez-Saldivar , G Villarreal-Villarreal . *Orthopaedics & Traumatology: Surgery & Research* 2017. 103  
146 (02) p. .

147 [Distal Radius Fracture: A Systematic Review of Observational Studies] *Distal Radius Fracture: A Systematic  
148 Review of Observational Studies*,

149 [Júnior et al. ()] ‘Epidemiologia e grau de satisfação do paciente vítima de trauma musculoesquelético atendido  
150 em hospital de emergência da rede pública brasileira’. Braga Júnior , MB , Chagas Neto , F A Porto , MA .  
151 *Acta Ortopédica Brasileira* 2005. 13 (3) p. .

152 [Fracture and Dislocation Classification Compendium Journal of Orthopaedic Trauma (2018)] ‘Fracture and  
153 Dislocation Classification Compendium’. *Journal of Orthopaedic Trauma* January 2018. Orthopaedic Trauma  
154 Association and AO Foundation. 32. (Supplement)

155 [Oliveira Filho et al. ()] ‘Fraturas do rádio distal: avaliação das classificações’. O M Oliveira Filho , W D  
156 Belangero , J B M Teles . *Revista da Associação Médica Brasileira* 2004. 50 (1) p. .

157 [Kruppa et al. ()] ‘Low complication rate of elastic stable intramedullary nailing (ESIN) of pediatric forearm  
158 fractures: A retrospective study of 202 cases’. C Kruppa , P Bunge , T A Schildhauer , M Dudda . *Medicine*  
159 2017. 96 (16) p. 6669.

160 [Collins ()] ‘Management and rehabilitation of distal radius fracture’. D C Collins . *Orthopedic Clinics of North  
161 America* 1993. 24 (2) p. .

162 [Zhang et al. ()] ‘Manual passive rehabilitation program for geriatric distal radius fractures’. W Zhang , L Wang  
163 , X Zhang , Q Zhang , B Liang , B Zhang . *Medicine* 2021. 100 (03) p. 24074.

164 [Slutsky and Herman ()] ‘Rehabilitation of distal radius fractures: a biomechanical guide’. D J Slutsky , M  
165 Herman . *Hand Clinics* 2005. 21 (3) p. .

166 [Ogliari et al. ()] ‘Seasonality of adult fragility fractures and association with weather: 12-year experience of a  
167 UK Fracture Liaison Service’. G Ogliari , T Ong , L Marshall , O Sahota . *Bone* 2021. 147 p. 115916.

168 [Reis et al. ()] ‘Terapia ocupacional no tratamento das fraturas do terço distal do rádio’. F B Reis , M C Corvelo  
169 , R M Françoso , F Faloppa , D Masieiro . *Folha Médica* 1990. 100 (1/2) p. .

170 [Johnson et al. ()] ‘The relationship between temperature and hip and wrist fracture incidence’. N A Johnson ,  
171 Erb Stirling , M Alexander , J J Dias . *The Annals of The Royal College of Surgeons of England* 2020. 102  
172 (5) p. .

173 [Albertoni et al. ()] ‘Tratamento das fraturas da extremidade distal do rádio’. W M Albertoni , F Faloppa , J C  
174 Belotti . *Revista Brasileira de Ortopedia* 2002. 37 (1/2) p. .