



Scan to know paper details and  
author's profile

# Nutritional Assessment of Advanced Cancer Patients Admitted to Home Palliative Care

*Sebastiano Mercadante, Giuseppe Massimo Bellavia, Flavio Fusco, Lucia Adamoli, Carla Scibilia, Alessio Lo Cascio & Alessandra Casuccio*

*University of Palermo*

## ABSTRACT

**Objectives:** The aim of this study was to investigate the nutrition profile of advanced cancer patients who were admitted to specialized home palliative care.

**Methods:** A consecutive sample of home care patients at admission of home care was assessed. Depression, anxiety, nausea, poor appetite, and poor well-being were measured by numerical scale 0-10. Mini nutritional assessment form (MNA-SF), fatigue assessment scale (FAS), and sarcopenia by SARC-F. were performed. The use of drugs used for anorexia, including corticosteroids, progestins, or others, was recorded.

**Results:** Data of 135 patients were analyzed. Sixty-eight percent and 77% of patients resulted to be malnourished and sarcopenic. In the multivariate regression analysis, anxiety ( $p= 0.036$ ) and total FAS ( $p=0.013$ ) were independently associated with malnutrition. Fifty-five percent of patients were receiving corticosteroids or megestol acetate. No significant associations with parameters examined were found.

**Keywords:** malnutrition, sarcopenia, advanced cancer, palliative care, home care.

**Classification:** NLM Code: WB 310

**Language:** English



Great Britain  
Journals Press

LJP Copyright ID: 392882

London Journal of Medical and Health Research

Volume 24 | Issue 4 | Compilation 1.0



© 2024. Sebastiano Mercadante, Giuseppe Massimo Bellavia, Flavio Fusco, Lucia Adamoli, Carla Scibilia, Alessio Lo Cascio & Alessandra Casuccio. This is a research/review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 4.0 Unported License <http://creativecommons.org/licenses/by-nc/4.0/>, permitting all noncommercial use, distribution, and reproduction in any medium, provided the original work is properly cited.



# Nutritional Assessment of Advanced Cancer Patients Admitted to Home Palliative Care

Sebastiano Mercadante<sup>a</sup>, Giuseppe Massimo Bellavia<sup>a</sup>, Flavio Fusco<sup>b</sup>, Lucia Adamoli<sup>c</sup>, Carla Scibilia<sup>a</sup>, Alessio Lo Cascio<sup>d</sup> & Alessandra Casuccio<sup>d</sup>

## ABSTRACT

**Objectives:** The aim of this study was to investigate the nutrition profile of advanced cancer patients who were admitted to specialized home palliative care.

**Methods:** A consecutive sample of home care patients at admission of home care was assessed. Depression, anxiety, nausea, poor appetite, and poor well-being were measured by numerical scale 0-10. Mini nutritional assessment form (MNA-SF), fatigue assessment scale (FAS), and sarcopenia by SARC-F. were performed. The use of drugs used for anorexia, including corticosteroids, progestins, or others, was recorded.

**Results:** Data of 135 patients were analyzed. Sixty-eight per cent and 77% of patients resulted to be malnourished and sarcopenic. In the multivariate regression analysis, anxiety ( $p=0.036$ ) and total FAS ( $p=0.013$ ) were independently associated with malnutrition. Fifty-five percent of patients were receiving corticosteroids or megestol acetate. No significant associations with parameters examined were found.

**Conclusion:** The majority of advanced cancer patients admitted to home palliative care were malnourished independently of the primary tumor diagnosis. Indeed, fatigue and anxiety were independently associated with malnutrition.

**Keywords:** malnutrition, sarcopenia, advanced cancer, palliative care, home care.

## Key Messages

1. Information regarding the nutritional pattern in advanced cancer patients admitted to specialized home care is lacking.

2. This study reported that a large number of patients admitted to specialized home palliative care are malnourished and sarcopenic.
3. This study suggests systematically assess the nutritional status and evaluate the need for an intervention according to expected survival.

**Author a σ cō** <sup>a</sup>: Regional home palliative care program, SAMOT, Palermo, Italy.

<sup>b</sup>: Palliative care unit, ASL3 Liguria, Genua, Italy.

<sup>a</sup> <sup>d</sup> <sup>x</sup>: Main Regional center for supportive and palliative care, La Maddalena Cancer center, Palermo, Italy @ University of Palermo.

## I. INTRODUCTION

About 18 million new cancer patients and 9.6 million cancer-related deaths occurred in 2018 worldwide. Many of these patients are living longer with metastatic disease due to advances in diagnostics and treatments [1]. Patients with severe, progressive, and incurable diseases, such as advanced cancer, should receive palliative care to improve quality of life through prevention and relief of suffering, pain, and other physical, psychosocial, and spiritual manifestations [2].

Palliative care in incurable cancer should be focused on symptom control, especially those impacting nutrition, and initiated simultaneously with active care at diagnosis since patients identify malnutrition as the primary cause of physical, cognitive, and social function loss [3]. It has been suggested that advanced cancer patients receiving palliative care should be assessed for nutritional deficiencies. Regardless of whether they are still on anticancer treatments or not, in these patients malnutrition leads to low performance status, impaired quality of life, unplanned hospitalizations, and reduced survival [4].

The role of nutritional support for cancer patients in palliative care is still a controversial issue [5]. In a recent review, the prevalence of weight loss and associated symptoms was quite heterogeneous. There was a large range of variability influenced by age and social factors, comorbidities, and setting of care. The advanced stage was found to be linked to a higher risk of developing malnutrition, as an expression of the relationship between tumor burden, inflammatory status, reduced caloric intake, and malabsorption [6].

Information regarding the nutritional status of advanced cancer patients admitted to home palliative care is lacking. Given the relevance of assessing nutritional and functional aspects of palliative care and the lack of studies addressing the topic, this study aimed to investigate the nutrition profile of advanced cancer patients who were admitted to specialized home palliative care.

## II. METHODS

This was a cross-sectional study of patients who were admitted to two specialized home palliative care program in Italy, in Palermo and Genua. A group of home care physicians adhered to the protocol and were trained to use of the tools selected for this study, with the help of a dietitian.

A consecutive sample of advanced cancer patients at admission of home care was assessed for a period of six months. Inclusion criteria were: age >18 years and advanced cancer disease. Exclusion criteria were: refusal of informed consent, expected survival less than one week, cognitive failure or incapacity to help assessment, and language barriers.

Epidemiological data was recorded, including age, gender, primary diagnosis, and Karnofsky level. Items of Edmonton Symptom Assessment Scale including depression, anxiety, nausea, poor appetite, and poor well-being were measured.

The following parameters were assessed: mini nutritional assessment form (MNA-SF), fatigue assessment scale (FAS), and the SARC-F. The MNA-SF is a screening tool to help identify

patients who are malnourished or at risk of malnutrition. It consists of six items: decline in food intake, weight loss, mobility, psychological stress or acute disease, neuropsychological problems, and the BMI. The scale is scored out of a total of 14 points and values below a threshold of 12 were used to identify at risk patients. The range 8-11 points corresponds to risk of malnutrition, and patients with 0-7 points are malnourished [7]. FAS is a validated tool, which has psychometric properties. Each item is scored on a 5-point scale from 1 (never) to 5 (always); total scores range from 10 to 50 [8]. We examined the presence/absence of clinically significant fatigue by dichotomizing the scale at the validated threshold: no fatigue (scores of 10-21) versus fatigue (scores of 22-50) [9]. SARC-F is a clinical index for sarcopenia screening. It includes five components: aid for walking, falls, getting up from a chair, climbing stairs, and strength. The scores for each item range from zero to two, obtaining a total score between zero to 10 points.

A score greater than or equal to four is defined as risk of sarcopenia [10]. Drugs used for anorexia, including corticosteroids, progestins, or others, were recorded, and when available, laboratory data. Patient's informed consent and local ethical committee approval were obtained.

### 2.1 Statistical Analysis

Absolute and relative frequencies have been reported for qualitative variables, and means and standard deviation (SD) for quantitative variables.

Frequency analysis was performed using the Pearson's chi-square test and Fisher exact test, as needed. An univariate and multivariable regression model was used to examine the correlations between malnutrition patient status (dependent variables), and clinical patient characteristics (independent variables). Odds ratios and related 95% confidence intervals (95% CI) were reported as well as the p-value. Data were analyzed by IBM SPSS Software 24 version (IBM Corp., Armonk, NY, USA). All p-values were two-sided and  $p \leq 0.05$  was considered statistically significant.

### III. RESULTS

Three-hundred-forty-three patients were consecutively assessed. Two-hundred-eight patients were not considered, according to the exclusion criteria: language barriers (n.8), no informed consent (n.13), poor prognosis-not collaborating (n.187).

Data of 135 patients were assessed. The principal characteristics of patients and parameters recorded are reported in Table 1.

Nine-two patients were malnourished and 35 were at risk of malnutrition (MNA-SF  $\leq 7$  and  $<12$ , respectively). One-hundred-four patients (77%) had sarcopenia (SARC-F $\geq 4$ ).

At univariate analysis, age ( $P=0.015$ ), Karnofsky ( $P=0.001$ ), depression ( $P=0.005$ ), anxiety ( $P=0.008$ ), poor appetite ( $P=0.031$ ), poor well-being ( $P=0.006$ ), total FAS ( $P=0.001$ ), total SARC-F ( $P<0.0005$ ), monocytes, ( $P=0.049$ ) eosinophils ( $P=0.047$ ), and d-dimer ( $P=0.045$ ) were significantly associated with malnutrition. In the multivariate regression analysis, anxiety (OR 2.849, 95% CI 1.070-7.591;  $p=0.036$ ) and total FAS (OR: 1.693 (95%CI 1.116-2.566;  $p=0.013$ ) were independently associated with malnutrition. Seventy-five patients were receiving corticosteroids or megestrol acetate. No significant associations with parameters examined were found.

*Table 1:* Principal characteristics and data of patients admitted to home care

Age (mean, SD)	75.6 (10.6)
Gender (m/f)	75/60
Karnofsky (mean, SD)	40.5 (10.2)
Primary tumor (n°)	135
Gastrointestinal	53
Lung	40
Urogenital	14
Hematologic	13
Breast	6
Head and neck	6
Other	3
Depression	4.3 (2.7)
Anxiety	4.3 (2.9)
Nausea	1.8 (2.6)
Poor appetite	4.7 (2.6)
Poor well-being	5.5 (2.4)
MNA-SF (mean, SD)	6.4 (3.0)
FAS (mean, SD)	29.7 (7.6)
SARC-F (mean, SD)	5.8 (2.6)

### IV. DISCUSSION

This study reported the data regarding the nutritional status of advanced cancer patients admitted to home care. About 68% of patients were malnourished, and 77% were sarcopenic.

There was an independent correlation between malnutrition and fatigue measured with FAS. Of

interest, a psychological symptom of ESAS, namely anxiety, was independently associated with malnutrition. While psychological symptoms invariably belong to the cluster of poor appetite, nausea, weakness, or poor-well-being [11-15], nutritional status has not been taken into consideration.

No differences among primary tumors was found. Pancreatic, esophageal, gastroenteric cancers, head and neck and lung cancer have been reported to be more likely associated with malnutrition [6]. This finding could be attributed to the very advanced stage of disease, flattening any difference among tumors, or may be the low number of patients in each tumor category.

Many patients were receiving drugs that are commonly used for a variety of symptoms other than improving appetite. The use of corticosteroids for the cluster of symptoms including, fatigue, lack of appetite, anorexia-cachexia, and well-being, remains controversial, as their efficacy may vary depending on the patients' general condition and seems to be reduced in patients with advanced stage of disease or poor performance status [16].

Corticosteroids and progestins have been found to have the potential to offer some benefit [17]. Recently, efficacy of corticosteroids for cancer-related cachexia was less relevant [16], than that reported in a previous randomized controlled trials [18,19]. Given the cross-sectional design of this study it is difficult to draw information regarding the effect of these drugs in a population with a limited survival at time of admission to home palliative care. On the other hand, it is widely accepted that aggressive feeding interventions are not appropriate at the end of life, and weight loss and reduced food intake are inevitable part of the dying process [20-24]. In a recent opinion expert paper, it has been reported that any form of clinical assisted nutrition is primarily indicated for the prevention of death from malnutrition in selected individuals from specific groups of patients with advanced cancer, for patients with an inability to introduce sufficient nutrients, and/or an inability to digest sufficient nutrients, but not indicated for the management of anorexia, weight loss, cancer cachexia, or reduced oral intake due to nutrition impact symptoms [25].

Data regarding nutritional assessment of advanced cancer patients admitted to home palliative care is poor, and prevalently based on the use of parenteral nutrition [26-28] or

performed at home with cancer and non-cancer diseases [29], or in other palliative care settings, namely inpatient units [30-32]. Malnutrition and poor quality of life are prevalent among advanced cancer patients in home palliative care. In a small study with a cross-sectional design in patients receiving home palliative care, 52.5% of patients were moderately or suspected of being malnourished and 32.8% of them were severely malnourished. Patient-Generated Subjective Global Assessment (PG-SGA) score used in this study showed a lower total quality of life, psychophysiological well-being, functional well-being and social/spiritual well-being [33]. In a cross-sectional multi-center Spanish study performed in different settings including oncology, radiotherapy, and home palliative care, more than 50% of patients had moderate-severe malnutrition, assessed by PG-SGA [34]. It is likely that in home palliative care patients malnutrition rate would be higher, although this distinction among the settings was not reported.

Patients with advanced cancer are commonly more malnourished compared to cancer patients in earlier stages [35]. In this study, where patients with poor immediate prognosis were excluded, survival was, however, relatively short, mostly in the range of 2-3 months, considering the large range of an advanced cancer status, which can be 6-24 months [4].

There are some limitations of this study, such as the large number of excluded patients. This limitation is frequently reported in studies of palliative care patients in the late stage of disease. Another limitation was the lack of monitoring of nutrient intake in these patients at this stage of disease for finding possible correlations with malnutrition, FAS, and SARC. Finally, the study lacks longitudinal assessment, having a cross-sectional evaluation at time of admission to home palliative care. However, this information may be useful for understanding the frequency of malnutrition in this population.

## V. CONCLUSION

The majority of advanced cancer patients admitted to home palliative care are

malnourished independently of the primary tumor diagnosis. Indeed, fatigue and anxiety were independently associated with malnutrition. Longitudinal studies may help to provide more information about these relationships. More studies should afford the question whether active treatments with orexant drugs or nutritional supplements may have a role in the comprehensive palliative care management to improve quality of life of very advanced cancer patients with a short survival time followed at home.

*No conflict of interest*

*No funding*

*Data available on request*

*Author contribution:*

SM: Conceptualization;  
 AL: Data curation; Supervision, writing original draft  
 AC: Formal analysis;  
 GB: Methodology  
 FF,GB, LA, CS ; Investigation;  
 All: Writing - review & editing.

## REFERENCES

- Bray, F.; Ferlay, J.; Soerjomataram, I.; Siegel, R.L.; Torre, L.A.; Jemal, A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J. Clin.* 2018, **68**, 394–424.
- World Health Organization (WHO), National cancer control programmes: policies and managerial guidelines. 2nd ed. Geneva: World Health Organization; 2002. <https://www.who.int/cancer/nccp/en/>.
- Lis CG, Gupta D, Lammersfeld CA, Markman M, Pankaj GV. Role of nutritional status in predicting quality of life outcomes in cancer e a systematic review of the epidemiological literature. *Nutr J* 2012; **11**: 27.
- Cotogni P, Stragliotto S, Ossola M, Collo A, Riso S. The Role of Nutritional Support for Cancer Patients in Palliative Care. *Cotogni Nutrients* 2021; **13**: 306.
- Thomas, K.; Wilson, J. A.; Gold Standards Framework. National Gold Standards Framework Centre in End of Life Care. The Gold Standards Framework Proactive Identification Guidance (PIG). The National GSF Centre's Guidance for Clinicians to Support Earlier Identification of Patients Nearing the End of Life, Leading to Improved Proactive Person-Centred Care. 2016. Available online: [www.goldstandardsframework.org.uk](http://www.goldstandardsframework.org.uk) (accessed on 5 December 2020).
- Bossi P, Delrio P, Mascheroni A, The Spectrum of Malnutrition/Cachexia/Sarcopenia in Oncology According to Different Cancer Types and Settings: A Narrative Review. *Nutrients* 2021; **13**: 1980.
- Rubenstein LZ, Harker JO, Salva A, Guigoz Y, Vellas B. Screening for undernutrition in geriatric practice: developing the short-form mini-nutritional assessment (MNA-SF). *J Gerontol A Biol Sci Med Sci* 2001; **56**: M366e72.
- Michielsen HJ, De Vries J, Van Heck GL. Psychometric qualities of a brief self-rated fatigue measure: the Fatigue Assessment Scale. *J Psychosom Res.* 2003; **54**: 345–52.
- Thong MSY, Mols F, Wang XS, Lemmens VEPP, Smilde TJ, van de Poll-Franse LV. Quantifying fatigue in (long-term) colorectal cancer survivors: a study from the population-based patient reported outcomes following initial treatment and long term evaluation of survivorship registry. *Eur J Cancer*. 2013; **49**: 1957–66.
- Malmstrom TK, Miller DK, Simonsick EM, Ferrucci L, Morley JE SARC-F: a symptom score to predict persons with sarcopenia at risk for poor functional outcomes. *Malmstrom TK, Miller DK, Simonsick EM, Ferrucci L, Morley JE. J Cachexia Sarcopenia Muscle.* 2016; **7**: 28–36.
- Nardi M, Catalini A, Galiano A, Santangelo OE, Pinto E, Feltrin A, Nucci D, Gianfredi V Association between malnutrition and anxiety in cancer patients-areetrospective study. *Support Care Cancer*. 2024; **32**: 152.
- Yennurajalingam S, Tayasanant S, Balachandran D, Padhye NS, Williams JL, Liu DD, Frisbee- Hume S, Bruera E Association

between daytime activity, fatigue, sleep, anxiety, depression, and symptom burden in advanced cancer patients: a preliminary report. *J Palliat Med* 2016; 19(8): 849–56.

13. Chabowski M, Polański J, Jankowska-Polańska B, Janczak D, Rosińczuk J. Is nutritional status associated with the level of anxiety, depression and pain in patients with lung cancer? *J Thorac Dis* 2018; 10(4): 2303–10.
14. Overcash J, Goetz R, Williams N. Prediction of Malnutrition and the Relationships Among Pain, Function, Depression, and Cancer Stage in Older Women With Breast Cancer. *Clin J Oncol Nurs.* 2023; 27: 411-7.
15. Gupta D, Lis CG, Granick J, et al. Malnutrition was associated with poor quality of life in colorectal cancer: a retrospective analysis. *J Clin Epidemiol*, 2006;59, 704-9.
16. Miura T, Okizaki A, Hasuo H, et al. Dexamethasone 8 mg for Cancer-Related Fatigue in Inpatients with Advanced Cancer Undergoing Palliative Care: A Multicenter Phase II Trial. *Palliat Med Rep* 2021; 2: 316-32.
17. Saeteaw M, Sanguanboonyaphong P, Yoodee J, et al. Efficacy and safety of pharmacological cachexia interventions: systematic review and network meta-analysis. *BMJ Support Palliat Care*. 2021; 11: 75-85.
18. Yennurajalingam S, Frisbee-Hume S, Palmer JL, et al.: Reduction of cancer related fatigue with dexamethasone: A double-blind, randomized, placebo-controlled trial in patients with advanced cancer. *J Clin Oncol* 2013; 31: 3076–82.
19. Paulsen Ø, Klepstad P, Rosland JH, et al.: Efficacy of methylprednisolone on pain, fatigue, and appetite loss in patients with advanced cancer using opioids: A randomized, placebo-controlled, double-blind trial. *J Clin Oncol* 2014; 32: 3221–8.
20. Cline D. Nutrition issues and tools for palliative care. *Home Healthc Nurse*. 2006; 24: 54-7.
21. Good P, Cavenagh J, Mather M, Ravenscroft P. Medically assisted nutrition for palliative care in adult patients. *Cochrane Database Syst Rev*. 2008; (4): Cd006274.
22. Arends J, Baracos V, Bertz H, et al. ESPEN expert group recommendations for action against cancer-related malnutrition. *Clin Nutr* 2017; 36: 1187-96.
23. Baxter JP, Fayers PM, Bozzetti F, et al. An international study of the quality of life of adult patients treated with home parenteral nutrition. *Clin Nutr*. 2019;38:1788-96.
24. Alley DE, Metter EJ, Griswold ME, et al. Changes in weight at the end of life: characterizing weight loss by time to death in a cohort study of older men. *Am J Epidemiol*. 2010; 172: 558-65.
25. Alderman B, Allan L, Amano K, et al. Multinational Association of Supportive Care in Cancer (MASCC) expert opinion/guidance on the use of clinically assisted nutrition in patients with advanced cancer. *Support Care Cancer* 2022; 30: 2983-92.
26. Cotogni P, Monge T, Passera R, Brossa L, De Francesco A. Clinical characteristics and predictive factors of survival of 761 cancer patients on home parenteral nutrition: A prospective, cohort study. *Cancer Med* 2020; 9: 4686-98.
27. Ruggeri E, Giannantonio M, Ostan R, Agostini F, Sasdelli AS, Valeriani L, Pironi L, Pannuti R. Choice of access route for artificial nutrition in cancer patients: 30 y of activity in a home palliative care setting. *Nutrition*. 2021; 90:111264.
28. Sowerbutts AM, Lal S, Sremanakova J, et al. Burden S. Palliative home parenteral nutrition in patients with ovarian cancer and malignant bowel obstruction: experiences of women and family caregivers. *BMC Palliat Care*. 2019; 18:120.
29. Van der Pols-Vijlbrief R, Wijnhoven HA, Molenaar H, Visser M. Factors associated with (risk of) undernutrition in community-dwelling older adults receiving home care: a cross-sectional study in the Netherlands. *Public Health Nutr*. 2016; 19: 2278-89.
30. Thoresen L, Fjeldstad I, Krogstad K, Kaasa S, Falkmer UG. Nutritional status of patients with advanced cancer: the value of using the subjective global assessment of nutritional status as a screening tool. *Palliat Med*. 2002; 16: 33-42.

31. Kwang AY, Kandiah M. Objective and subjective nutritional assessment of patients with cancer in palliative care. *Am J Hospice Palliative Med.* 2010; 27: 117-26.
32. Stevens T, Keller H, Williams N, Downar J, Guthrie DM. Cross-Sectional Nutrition Profile of Palliative Home Care Clients in Ontario and Performance of the interRAI Palliative Care Nutrition Clinical Assessment Protocol. *JPEN J Parenter Enteral Nutr* 2021;45:183-192.
33. Shahmoradi N, Kandiah M, Peng LS. Impact of nutritional status on the quality of life of advanced cancer patients in hospice home care *Asian Pac J Cancer Prev* 2009; 10: 1003 -09.
34. Segura A, Pardo J, Jara C, Azuara ML, Casado J. An epidemiological evaluation of the prevalence of malnutrition in Spanish patients with locally advanced or metastatic cancer. *Clin Nutr* 2005;24:801-14.
35. Ravasco P, Monteiro-Grillo I, Vidal P, et al. Nutritional deterioration in cancer: the role of disease and diet. *Clin Oncol*, 2003;15, 443-50.

*This page is intentionally left blank*